

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/311845401>

Transference-focused psychotherapy reduces treatment drop-out and suicide attempters compared with community psychotherapist treatment in borderline personality disorder

Article in *Evidence-Based Mental Health* · October 2011

DOI: 10.1136/ebmh1097

CITATIONS

6

READS

196

3 authors, including:



Kenneth Levy

Pennsylvania State University

219 PUBLICATIONS 10,515 CITATIONS

[SEE PROFILE](#)



Frank E. Yeomans

Weill Cornell Medical College

97 PUBLICATIONS 2,462 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Interpersonal Match as a Predictor of Marital Satisfaction and PTSD in Military Couples [View project](#)



Tend and Befriend & Attachment [View project](#)

Transference-focused psychotherapy reduces treatment drop-out and suicide attempters compared with community psychotherapist treatment in borderline personality disorder

QUESTION

Question: How does transference-focused psychotherapy (TFP) compare with treatment by community psychotherapists for borderline personality disorder (BPD)?

Patients: One hundred and four female outpatients aged between 18 and 45 years with Diagnostic and Statistical Manual, Fourth Edition (DSM-IV) BPD. Main exclusions: anti-social personality disorder, schizophrenia, bipolar I or II disorder with an episode in the past 6 months, recent substance dependency, organic pathology or mental retardation.

Setting: Two outpatient units, Germany; recruitment October 2004 to August 2006.

Intervention: TFP or community psychotherapist treatment for a year. TFP was a manualised-modified psychodynamic psychotherapy delivered in two 50 min sessions per week by a specially trained therapist. Community psychotherapists were experienced and particularly interested in people with BPDs; however, none of the therapists had specific training in manualised borderline psychotherapy.

Outcomes: (1) Primary outcomes included drop-out from treatment, suicide attempts (assessed using the Cornell Interview for Suicidal and Self Harming Behaviour – Self Report). (2) Secondary outcomes included DSM-IV BPD diagnosis. Participants were assessed at baseline and 1 year.

Patient follow-up: Sixty-six per cent completed treatment (see Results); 69% completed final assessment and were included in observed case (OC) analysis; 100% included in last observation carried forward (LOCF) analysis.

METHODS

Design: Randomised controlled trial.

Allocation: Not clear.

Blinding: Single blind (outcome assessors blinded).

Follow-up period: One year (treatment period only).

MAIN RESULTS

Drop-out from treatment was significantly higher in the community psychotherapy group than in the TFP group (38.5% with TFP vs 67.3% with community treatment; $p=0.003$). TFP reduced the proportion of participants making suicide attempts compared with community psychotherapist treatment (AR of a suicide attempt, LOCF analysis: 13.7% with TFP vs 21.2% with community psychotherapy; $p=0.009$; OC analysis ARs not presented, $p=0.001$). After 1 year, TFP reduced the proportion of participants who met DSM-IV criteria for BPD compared with community psychotherapist treatment (AR for not meeting DSM-IV criteria for BPD, OC analysis: 51.2% with TFP vs 27.6% with community psychotherapy, $p=0.047$; LOCF analysis: 42.3% with TFP vs 15.4% with community psychotherapy, $p=0.002$).

CONCLUSIONS

Among women with BPD, TFP reduces treatment drop-out and the proportion of people making suicide attempts compared with treatment by a community psychotherapist. TFP also reduces the proportion of women continuing to meet diagnostic criteria for BPD at 1 year.

ABSTRACTED FROM

Doering S, Hörz S, Rentrop M, et al. Transference-focused psychotherapy v. treatment by community psychotherapists for borderline personality disorder: randomised controlled trial. *Br J Psychiatry* 2010;**196**:389–95.

Correspondence to: Stephan Doering, Psychosomatics in Dentistry, Department of Prosthodontics and Material Sciences, University of Muenster, Waldeyerstrasse 30, 48149 Muenster, Germany; stephen.doering@ukmuenster.de

Source of funding: Austrian National Bank.

Additional references are available online only at <http://ebmh.bmj.com>

Borderline personality disorder (BPD) is a highly prevalent and debilitating disorder (1–6% in the general population and 9–15% in outpatient samples). BPD is frequently comorbid with a number of Axis I disorders, when comorbid BPD negatively affects the course and outcome of otherwise efficacious treatments for these disorders.¹ Thus, the treatment of BPD is a major public health concern.

The randomised controlled trial by Doering and colleagues further establishes a psychodynamic treatment called transference-focused psychotherapy (TFP) as an efficacious treatment for BPD. The authors compared 1 year of TFP to treatment by experienced community psychotherapists (ECP). While patients improved in both treatments, patients randomly assigned to TFP evidenced significantly lower drop-out, greater reductions in patients attempting suicide, lower number of inpatient admissions, fewer patients experienced BPD symptoms and greater improvements in personality organisation and psychosocial functioning.

Strengths of this study include (1) a large sample with the power to detect differences between the groups; (2) reporting of both completer (internal validity) and intent-to-treat analyses (external validity); (3) comparisons with therapists in the community who were both experienced and committed to treating BPD patients (ecological validity/legitimate control group); and (4) the demonstration of efficacy for TFP by an independent group not affiliated with the original investigators.

One limitation is that those in the TFP condition received significantly more psychotherapy sessions than in the ECP condition (2:1 for completers; 3:1 for Intention to Treat). However, the authors statistically controlled for the sessions and found no effect on the treatment outcome. Additionally, in most other BPD studies the experimental group tends to have more treatment hours than the control (with 60:1², 6:1³ and 2:1⁴ in prior studies).

These findings contribute to a growing literature on efficacious approaches available to clinicians treating BPD. There is no credible evidence that any

one treatment is significantly better than any other as a function of effect sizes or comparisons with bona fide alternative treatments. Thus, TFP represents one of a number of therapies that may be useful in treating BPD. Additionally, only TFP has been shown to change how patients think about themselves in relationships.⁵ Future research is needed on the long-term efficacy of these approaches, identifying mechanisms of change and determining which patients will do best in which specific treatments.

Kenneth N Levy,^{1,3} Kevin B Meehan,² Frank E Yeomans³

¹Department of Psychology, Pennsylvania State University, University Park, Pennsylvania, USA

²Department of Psychology, Long Island University, Brooklyn, New York, USA

³Department of Psychiatry, Joan and Sanford I. Weill Medical College, Cornell University, New York, USA

Competing interests None.

EBMH

Transference-focused psychotherapy reduces treatment drop-out and suicide attempters compared with community psychotherapist treatment in borderline personality disorder

Evid Based Mental Health 2010 13: 119 originally published online August 10, 2010
doi: 10.1136/ebmh1097

Updated information and services can be found at:
<http://ebmh.bmj.com/content/13/4/119>

References

These include:

This article cites 6 articles, 1 of which you can access for free at:
<http://ebmh.bmj.com/content/13/4/119#BIBL>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Suicide \(psychiatry\)](#) (228)
[Borderline personality disorder](#) (33)
[Clinical trials \(epidemiology\)](#) (989)
[Epidemiology](#) (1570)
[Antisocial personality disorder](#) (37)
[Neurology](#) (1070)
[Schizophrenia spectrum](#) (430)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>